

# Suicide Prevention in Youth and Young Adults

## Communicating With Families Saves Lives

A Checklist for Health Providers  
and Mental Health Practitioners

Created by the *Oregon Council of  
Child and Adolescent Psychiatry*



### Rationale and Scope:

Oregon's suicide rate in 2010, at 15.2 per 100,000 persons, was 35% above the national average. The rate in 2010 among Oregon males ages 20-24 was 22.6 per 100,000. Suicide ranks as the second leading cause of death in Oregon among youth ages 10-24. Suicide claims more lives each year nationally than homicide, HIV/AIDS, or auto accidents. On average, 45% of all suicide victims nationally had contact with primary care providers within one month of taking their own lives.

Communication between primary care providers and/or mental health practitioners and family members of patients seeking treatment for mental illness improves the quality of care provided to these patients, reduces the risk of suicide and self-harm behaviors, and encourages the use of resources to improve overall outcomes for these patients. While confidentiality is a fundamental component of a therapeutic relationship, it is not an absolute, and the safety of the patient overrides the duty of confidentiality. Misunderstandings by clinicians about the limitations created by HIPAA, FERPA, and state laws for preserving confidentiality of patients has caused unnecessary concern regarding disclosure of relevant clinical information. Communication between providers, patients, and family members/identified significant others needs to be recognized as a clinical best practice and deviations from this should occur only in rare and special circumstances.

This checklist is intended for use by primary care providers, emergency department staff, and any professional providing mental health treatment, to include, but not be limited to, family physicians, general practitioners, pediatricians, physician assistants, nurse practitioners, social workers, counselors, psychologists, psychiatric nurse practitioners and psychiatrists. If your professional organization does not have a preferred suicide risk assessment protocol, please see page 7 of this document.

Patient Name: \_\_\_\_\_

Date: \_\_\_\_\_

DOB/Medical Record #: \_\_\_\_\_

**For All Patients with Mental Health Issues, Providers Should Complete the Following:**

- Complete a comprehensive risk assessment including patient interview, record review and communication with family/parents. **If you do not feel qualified to complete a comprehensive risk assessment, refer the patient for urgent evaluation and verify completion.**
  - If the patient is 18 or older, or if you believe confidentiality is required by law or common medical practice, seek an authorization to release information for the family/parents **or** document a compelling reason not to do so. Be assertive and persuasive in obtaining this authorization.
  - Interview the family, to the extent they are involved in the patient's care, to obtain additional history about the patient and to determine what the family/parents already know about the illness/need for treatment.
  - Obtain authorizations to release information for all previous treatment providers and promptly request treatment records, including psychotherapy, psychiatric and relevant medical records.
  - Review the medical records carefully to gain a comprehensive knowledge of risk factors for the patient.

**Where an elevated risk of suicide is identified or patient is a minor in Oregon (see notes, page 4):**

- Following the initial evaluation, communicate with the patient and the family/parents regarding diagnoses, treatment recommendations and safety issues. Do **not** assume they know anything about the nature of mental illness, treatment, risk factors, or community resources.
- Regardless of whether one has a signed authorization, explicitly inform the family in the presence of the patient of all safety issues, including risk factors for suicide and what steps to take if danger exists, such as ridding the home of firearms/other means of self-harm and creating a plan to monitor and support the patient.
- Regardless of whether one has a signed authorization, discuss available community resources to help the family and patient, including resources for case management, support groups, improving mental health at home, and other relevant factors.
- Coordinate provision of care when a patient transitions from one level of care to another, or one provider to another:
  - Involve patient and family in planning process including discussion of interim safety plan.
  - Assure follow up is in place with a specific timely appointment.
  - Assure accepting provider has full knowledge of history and risk issues/records.
  - Confirm that patient has attended the follow up appointment.

**Additional Notes:**

Clinician Signature: \_\_\_\_\_

Date: \_\_\_\_\_

## **Sources and Justifications:**

### **Purpose:**

The basic philosophy behind this document is to always do what is in the best interest of the patient in the provider's professional judgment. OCCAP believes that in nearly all cases, this will involve a great deal of communication with the families of patients. We also believe that the perception of restrictions on communication with families is often greater than the limits imposed by law or regulation.

This checklist is created with the needs of children, adolescents, and transition-age youths (young people between the ages of 17 and 24) in mind. (It may of course be used beneficially with older patients as well.) It is widely recognized that for these patients the best outcomes require the provider(s) to communicate with the family and involve them in treatment.

A common misperception is that this approach is prohibited by law or ethical standards. The most obvious recourse is to seek authorization for release of information, and advocate with the patient to sign the authorization at an early stage in treatment (APA April 1999). However, both legal and ethical standards provide for communication with parents/family under many circumstances even without the patient's authorization.

A patient seeking care from an abusive family situation or youth seeking care only on the condition of confidentiality must be treated as special cases, and the discretion of the treatment provider should be exercised. We believe that for the vast majority of treatment scenarios, however, involvement of the family will enhance care and reduce both the risk of suicide and the risk of a malpractice suit.

"From a liability risk standpoint, suicide claims are considered to be low frequency/high severity. The overall number of suicide-related malpractice cases is low; however, those that are filed tend to result in higher than average indemnity payments. The issue in these cases is NOT whether the physician should have been able to predict the patient would commit suicide, but, rather: was the risk of suicide foreseeable? Physicians who actively address suicide risk, and document their risk assessment and recommended treatment plan, are far less likely to be named in a lawsuit alleging improper care resulting in suicide." (Debra Davidson, American Physicians Assurance Corporation)

### **Key Points:**

1. Always act in the best interests of your patient, with particular attention to the safety of that individual. **If you are in doubt, it is better to violate confidentiality in the interest of protecting your patient.**
2. It is **always** better to defend allegations of an inappropriate disclosure than to defend a failure to disclose with bad outcome (e.g. injury or death).
3. Do not assume that confidentiality is required or that a release is necessary to communicate with parents, if it cannot be obtained. State law might indicate otherwise. (See ORS 109.650, 109.680 and 109.675 on when a provider may or must disclose to parents.)
4. Where you believe that confidentiality may be required and you cannot obtain a release, communicate with the family/parents non-confidential information such as community resources available to them to help address the issue. For example, if the family knows that the patient is clinically depressed and that there has been non-suicidal self-injury (such as cutting), give the family/parents general information on depression and non-suicidal self-injury, and on community resources where they may obtain more information and support.
5. Approach safety planning and treatment as a collaborative process from the onset. Assure the patient that s/he is not alone; engage their identified support system. Recognize that family may provide information to you without the need for a release. Consider also non-traditional sources of information such first responders, police, ER staff.

## **Laws, Ethics, and Interpretations:**

It may be that **state law** allows or even requires the provider to involve the family at some stage of the treatment, when the disclosure or involvement is clinically appropriate. (See ORS 109.650, 109.680 and ORS 109.675 [2] on when the provider may or must disclose.)

### **Oregon Laws Regarding Disclosure of Protected Health Information:**

#### **109.650 Disclosure without minor's consent and without liability.**

A hospital or any physician, nurse practitioner or dentist as described in ORS 109.640 may advise the parent or parents or legal guardian of any minor of the care, diagnosis or treatment or the need for any treatment, without the consent of the patient, and any hospital, physician, nurse practitioner or dentist is not liable for advising the parent, parents or legal guardian without the consent of the patient. [1971 c.381 §2; 2005 c.471 §8]

#### **109.675 Right to diagnosis or treatment for mental or emotional disorder or chemical dependency without parental consent.**

(1) A minor 14 years of age or older may obtain, without parental knowledge or consent, outpatient diagnosis or treatment of a mental or emotional disorder or a chemical dependency, excluding methadone maintenance, by a physician licensed by the Oregon Medical Board, a psychologist licensed by the State Board of Psychologist Examiners, a nurse practitioner registered by the Oregon State Board of Nursing, a clinical social worker licensed by the State Board of Licensed Social Workers or a community mental health program established and operated pursuant to ORS 430.620 when approved to do so by the Oregon Health Authority pursuant to rule.

(2) *However, the person providing treatment shall have the parents of the minor involved before the end of treatment unless the parents refuse or unless there are clear clinical indications to the contrary, which shall be documented in the treatment record.* (Emphasis added) The provisions of this subsection do not apply to:

(a) A minor who has been sexually abused by a parent; or

(b) An emancipated minor, whether emancipated under the provisions of ORS 109.510 and 109.520 or 419B.550 to 419B.558 or, for the purpose of this section only, emancipated by virtue of having lived apart from the parents or legal guardian while being self-sustaining for a period of 90 days prior to obtaining treatment as provided by this section. [1985 c.525 §1; 1989 c.721 §47; 1993 c.546 §137; 1997 c.249 §38; 2009 c.442 §30; 2009 c.595 §71]

See also, Oregon Health Authority: Standards for the Approval of Providers of Non-Inpatient Mental Health Treatment Services implemented pursuant to ORS 109.695.

OAR 309-039-0540 General Standards: Each provider is required to meet the following administrative standards . . . (12)(a)“Providers shall provide for the **earliest feasible involvement** of the parents or guardians in the treatment plan consistent with the clinical requirement of the minor as provided under ORS 109.695.” (Emphasis added.)

#### **109.680 Disclosure without minor's consent; civil immunity.**

A physician, psychologist, nurse practitioner, clinical social worker licensed under ORS 675.530 or community mental health program described in ORS 109.675 may advise the parent or parents or legal guardian of any minor described in ORS 109.675 of the diagnosis or treatment whenever the disclosure is clinically appropriate and will serve the best interests of the minor's treatment because the minor's condition has deteriorated or the risk of a suicide attempt has become such that inpatient treatment is necessary, or the minor's condition requires detoxification in a residential or acute care facility. If such disclosure is made, the physician, psychologist, nurse practitioner, clinical social worker licensed under ORS 675.530 or community mental health program shall not be subject to any civil liability for advising the parent, parents or legal guardian without the consent of the minor. [1985 c.525 §2; 1989 c.721 §48; 2009 c.442 §31; 2009 c.595 §72]

See also ORS 441.750 (authorizing certain disclosures and prohibiting civil or criminal liability provided physician is acting in good faith.)

**Federal law** permits, but does not require, a provider to disclose to family protected health information relevant to the family member's involvement in the patient's care without a release if the provider provides an opportunity for the patient to object and the patient does not, or the provider "reasonably infers from the circumstances, based [on] the exercise of professional judgment, that the [patient] does not object to the disclosure." (45 CFR 164.510 [b][2])

Providers may also disclose otherwise protected information when "the opportunity to agree or object to the use or disclosure cannot practicably be provided because of the individual's incapacity or an emergency circumstance, the [provider] may, in the exercise of professional judgment, determine whether the disclosure is in the best interests of the individual . . ." (45 CFR 164.510 [b][3])

"Medical professionals can talk freely to family and friends, unless the patient objects. No signed authorization is necessary. Health care workers may not reveal confidential information about a patient or case to reporters, but they can discuss general health issues." Susan McAndrew, Deputy Director of Health Information Privacy, U.S. Department of Health and Human Services, quoted in: Gross, Jane. *Keeping Patients' details private, even from kin*. New York Times, July 3, 2007.

Also note, if a parent seeks access to medical records of an un-emancipated minor, federal law (HIPAA) defers to any state law which specifically permits or prohibits disclosure, e.g. Oregon statutes. (45 CFR 164.502 (g)(3)(ii); see also State of Oregon, Legislative Counsel Committee, *Disclosure of protected health information to a parent*, 10/18/2011, and OMA Legal/Regulatory Briefing on Specially Protected Records, page 5.)

Justifiable exceptions exist even when one of these laws does not apply. The **American Psychiatric Association** has said "Disclosures can sometimes be justified on the grounds that they are necessary to protect the patient. For instance, it is generally acceptable for a psychiatrist to warn a patient's family or roommate when the patient is very depressed and has voiced suicidal thoughts." (Practice Management Handbook for Early Career Psychiatrists, Office of Healthcare Systems and Financing, American Psychiatric Association, June 2010; also see 34 CFR 99.36 as regards educational institutions.)

Specifically related to college students, the **American Psychiatric Association** has said, "Parents also have a strong interest in being involved in their children's health care—even when their child might legally be an adult... The perceived impediments to disclosures by college officials in situations in which the health or safety of students may be endangered seem often to be the result of a misunderstanding of FERPA and other relevant laws and regulations." The paper notes that suicide is the second leading cause of death among American college students, and the third leading cause among all American youth, aged 10-24. (American Psychiatric Association, *College Mental Health and Confidentiality*, June 2009)

The **Oregon Medical Association** has said as regards communication with other providers, that "It is important to keep in mind that, while it is sound practice to evaluate what information needs to be disclosed, it is also important not to be too restrictive. If the appropriate information is not disclosed for treatment purposes, it can lead to adverse consequences for the patient... Some of the more restrictive practices that have been adopted by some physicians represent a barrier to efficient and quality health care." (Oregon Medical Association HIPAA Privacy Rule Overview, accessed through <http://www.theoma.org/practice-help-center/practice-management/privacy-security/hipaa/hipaa-privacy-rule-overview> on 9/16/2011)

The **American Association of Suicidology** (AAS) has prepared "Recommendations for Inpatient and Residential Patients Known to be at Elevated Risk for Suicide". Recommendation number 7 states, "Both the patient and the family or significant others should be given instruction regarding suicide and its associated risk, including, but not limited to the following: warning signs of suicide, the increased risk for suicide during pass or following discharge; the need for medication and other treatment adherence; explanation of how psychiatric symptoms may impair judgment; explanation of the need for the patient to avoid use of intoxicants and how

intoxicants increase risk; the need for the removal of the means for suicide, and the particular risk associated with firearms.” This reasoning is equally valid to out-patient care.

([http://www.suicidology.org/c/document\\_library/get\\_file?folderId=231&name=DLFE-106.pdf](http://www.suicidology.org/c/document_library/get_file?folderId=231&name=DLFE-106.pdf))

Also, it is important to note that one cannot breach confidentiality where it does not exist, or where one is not disclosing medical records or information learned from the patient. For instance, if the family brings you prior treatment records, you may discuss the prior treatment with them and its implications for the current situation plus provide referral to community support such as NAMI.

**Generally accepted ethical standards** also promote greater communication with families and/or significant others.

### **American Academy of Child and Adolescent Psychiatry Practice Parameter on Depressive Disorders (2007):**

**Recommendation 1.** The Clinician Should Maintain a Confidential Relationship With the Child or Adolescent While Developing Collaborative Relationships With Parents, Medical Providers, Other Mental Health Professionals, and Appropriate School Personnel [MS\*].

At the outset of the initial contact, the clinician should clarify with the patient and parents the boundaries of the confidential relationship that will be provided. The child’s right to a confidential relationship is determined by law that varies by state. Each state has mandatory child abuse reporting requirements. Parents will expect information about the treatment plan, the safety plan, and progress toward goals of treatment. The child should expect that suicide or violence risk issues will be communicated to the parents. (*Emphasis added*). The clinician should request permission to communicate with medical providers, other mental health professionals involved in the treatment, and appropriate school personnel. Clinicians should provide a mechanism for parents to communicate concerns about deterioration in function and high-risk behaviors such as suicide threats or substance use.

**Recommendation 7.** Each Phase of Treatment Should Include Psychoeducation, Supportive Management, and Family and School Involvement [MS\*]...

*Family Involvement.* Even in the absence of formal family therapy, it is virtually impossible to successfully treat a child or adolescent patient without the close involvement of parents. (*Emphasis added*). First, the clinician has to recognize that motivation for treatment comes often from the parents, and therefore the treatment contract must involve them. Second, the parents may observe aspects of the child’s functioning or symptoms that the child either is not aware of or does not wish to share, and this information is vital to the development of a realistic and effective treatment contract. Third, the parents are able to monitor their child’s progress and serve as a safety net. As described in the section about psychotherapies (Recommendation 9), despite the scarce and weak empirical evidence, knowledge of risk factors suggests that interventions with families are an important part of clinical management (*Emphasis added*). These interventions should take into account the family’s cultural and religious background and focus on strengthening the relationship between the identified patient and caregiver(s), provide parenting guidance (e.g., management of conflicts), reduce family dysfunction, and facilitate treatment referral for caregivers or siblings with psychiatric disorders and for marital conflict (Asarnow et al., 1993 [rct]; Birmaher et al., 2000 [ut]; Diamond et al., 2002 [ut]; Garber et al., 2002; Hammen et al., 2004; Nomura et al., 2002; Sanford et al., 2006)...

### **American Psychiatric Association Practice Guideline for the Assessment and Treatment of Patients With Suicidal Behaviors (2003):**

Section V: Documentation and Risk Management, *C. Communication With Significant Others*

The confidential nature of the doctor-patient relationship is a fundamental aspect of the psychotherapeutic process. Consequently, the psychiatrist will need to manage the tension between this requirement and the wish to act in the patient’s best interest. (*Emphasis added*). The default position is to maintain confidentiality unless the patient gives consent to a specific intervention or communication. However, in maintaining a safe environment for the patient, significant others may need to be contacted to furnish historical information or carry out specific tasks such as removing firearms from the home. If the psychiatrist determines that the patient is (or is likely to become) dangerous to him- or herself or to others and the patient



will not consent to interventions that aim to reduce those risks, then the psychiatrist is justified in attenuating confidentiality to the extent needed to address the safety of the patient and others (*Emphasis added*). More specifically, the 2001 edition of [The American Medical Association] *The Principles of Medical Ethics With Annotations Especially Applicable to Psychiatry* states: "[P]sychiatrists at times may find it necessary, in order to protect the patient or the community from imminent danger, to reveal confidential information disclosed by the patient" (Section IV, Annotation 8). As with many situations involving the suicidal patient, such decisions require much clinical judgment in weighing the effects of breaching confidentiality on the therapeutic relationship against the potential safety risks for the patient or others. It should also be noted that the psychiatrist can listen to information provided by friends or family without violating confidentiality by disclosing information about the patient to the informant. In addition, in an emergency situation, necessary information about the patient can be communicated with police and with emergency personnel, including medical staff and emergency medical technicians.

\* The AACAP Practice Parameters specify the strength of the underlying clinical and/or empirical data of the recommendations; the [MS] designation reflects a "Minimal Standard" described in the practice parameters as: "Minimal Standards are applied to recommendations that are based on rigorous empirical evidence (e.g., randomized controlled trials) and/or overwhelming clinical consensus. Minimal standards apply more than 95% of the time (i.e., in almost all cases)."

### **Suicide Risk Assessment Resources for Clinicians:**

1. SAFE-T (Suicide Assessment Five-step Evaluation and Triage) available to download at [www.sprc.org](http://www.sprc.org). Go to the library and resources section, browse the library, search for SAFE-T, or copy and paste this: [http://www.sprc.org/sites/sprc.org/files/library/safe\\_t\\_pcktrd\\_edc.pdf](http://www.sprc.org/sites/sprc.org/files/library/safe_t_pcktrd_edc.pdf)
2. An excellent screening resource developed specifically for PCPs can be found at: <http://prevention.mt.gov/suicideprevention/PrimaryCarePhysiciansMasterFile.pdf>. The Master File has many other easy to use tools on suicide prevention and is simple to access.

### **References:**

- American Academy of Child and Adolescent Psychiatry, *Practice Parameter on Depressive Disorders*, (2007)
- American Psychiatric Association, *Practice Guideline for the Assessment and Treatment of Patients with Suicidal Behavior*, 2003.
- American Psychiatric Association, *College Mental Health and Confidentiality Resource Document*, 2009.
- American Psychiatric Association, *Office of Health Care Systems and Financing, Chap. 20, Confidentiality*, June 2010.
- Appleby, L., Shaw, J., Amos, T., McDonnell, R., Harris, C., McCann, K., et al. (1999). Suicide within 12 months of contact with mental health services: national clinical survey. *British Medical Journal*, 318, 1235-1239.
- Busch, K. A., Fawcett, J., & Jacobs, D. G. (2003) Clinical correlates of inpatient suicide. *Journal of Clinical Psychiatry*. 64 (1), 14-19.
- Ho, T.P. (2003) The suicide risk of discharged psychiatric patients. *Journal of Clinical Psychiatry*, 64 (6), 702-707.
- Luoma, J. B., Pearson, J. L., & Martin, C. E. (2002). Contact with Mental Health and Primary Care Prior to Suicide: A Review of the Evidence. *American Journal of Psychiatry*, 159, 909-916.
- State of Oregon, Legislative Counsel Committee, *Disclosure of protected health information to a parent*, 10/18/2011.
- Oregon Medical Association, *Regulatory Briefing: Specially Protected Records*, 10/1/2010.
- Oregon Medical Association, *HIPAA Privacy Rule Overview*, 6/14/2011.
- Office of the Ombudsman for Mental Health and Mental Retardation, State of Minnesota, *Suicide Prevention Alert*, February, 2002.
- Suicide Prevention Resource Center, *Continuity of Care for Suicide Prevention and Research*, 2011.

**Publication Data:**

This document was adopted as the official publication and position of the Oregon Council of Child and Adolescent Psychiatry on June 5, 2012 and authorized for unlimited distribution and use. The checklist itself is NON-PROPRIETARY and may be edited and revised to be specifically applicable to any clinical setting, as long as the edits are clearly indicated as not being part of the original document.

Principal authors are Stewart S. Newman, MD and G. Ness Matthew Gabay, JD. Ajit Jetmalani, MD and Kirk Wolfe, MD provided significant editing and consultation. We are grateful to the numerous attorneys, physicians, medical faculty, advocates and mental health practitioners who reviewed the document and made helpful suggestions. The Executive Committee of OCCAP reviewed the document twice in the course of a year during its development. The authors have no relevant potential conflicts of interest to disclose regarding this project.

The checklist and resource material are available for downloading and printing on the Internet at [www.aacap.org/occap](http://www.aacap.org/occap) or [www.occap.org](http://www.occap.org) as Communication Checklist for Mental Health Practitioners. Requests for reprints may be made to OCCAP, PO Box 2042, Salem, OR 97308.

**Legal Disclaimer: The material in this document has been prepared for informational purposes only and does not constitute legal advice.**